arXiv:1609.06573v1 [math.GT] 21 Sep 2016

UNKNOTTING SUBMANIFOLDS OF THE 3-SPHERE BY TWISTINGS

MAKOTO OZAWA

Abstract. By the Fox's re-embedding theorem, any compact submanifold of the 3-sphere can be re-embedded in the 3-sphere so that it is unknotted. It is unknown whether the Fox's re-embedding can be replaced with twistings. In this paper, we will show that any closed 2-manifold embedded in the 3-sphere can be unknotted by twistings. In spite of this phenomenon, we show that there exists a compact 3-submanifold of the 3-sphere which cannot be unknotted by twistings. This shows that the Fox's re-embedding cannot always be replaced with twistings.

1. Introduction

Throughout this paper, we will work in the piecewise linear category. We assume that a surface is a compact, connected 2-manifold and that a 2-manifold is possibly disconnected.

Definition 1.1. Let X be a compact submanifold of the 3-sphere S^3. Take a loop C in $S^3 - X$ which is the trivial knot in S^3. Then C bounds a disk D in S^3, which may intersect X in its interior. Cut open S^3 along by D, rotate one copy of D by $\pm 2\pi$, and glue again two copies of D. Then we obtain another submanifold X' of S^3 and call this operation a *twisting along* C, which is denoted by $(S^3, X) \xrightarrow{\mathrm{C}} (S^3, X')$.

Remark 1.2. A twisting along C is not a homeomorphism of S^3, but it gives a homeomorphism of $S^3 - C$. We note that a twisting along C is also obtained by ± 1-Dehn surgery along C.

Definition 1.3. Let X be a compact submanifold of S^3 which has an n connected components $X_1, \ldots, X_n$.

We say that $X = X_1 \cup \cdots \cup X_n$ is *completely splittable* in S^3 if there exist $n-1$ mutually disjoint 2-spheres $S_1, \ldots, S_{n-1}$ in $S^3 - X$ such that if we cut open S^3 along $S_1 \cup \cdots \cup S_{n-1}$ and glue $2(n-1)$ 3-balls along their boundaries, then we obtain n pairs of the 3-sphere and the submanifold $(S^3, X_1), \ldots, (S^3, X_n)$.

For a connected component X_i of X, we say that a pair (S^3, X_i) is *unknotted* in S^3 if the exterior $E(X_i) = S^3 - intN(X_i)$ consists of handlebodies. We say that X is *unknotted* if X is completely splittable and for every pair (S^3, X_i), X_i is unknotted in S^3.

2010 *Mathematics Subject Classification.* 57Q35 (Primary), 57N35 (Secondary).

Key words and phrases. embedding, closed surface, 3-sphere, twisting, crossing changes, unknotting, Fox's re-embedding.

The author is partially supported by Grant-in-Aid for Scientific Research (C) (No. 26400097), The Ministry of Education, Culture, Sports, Science and Technology, Japan.

Remark 1.4. We remark that by the Fox's re-embedding theorem [2], any compact submanifold M of S^3 can be re-embedded in S^3 so that M is unknotted.

The following is the main subject of this paper.

Problem 1.5. *Can any Fox's re-embedding be replaced with twistings?*

It is well-known that Problem 1.5 is true for any closed 1-manifold and for any closed 2-manifold which bounds handlebodies.

Definition 1.6. Let F be a closed 2-manifold and α be a loop, namely, simple closed curve in F. We say that α is *inessential* in F if it bounds a disk in F. Otherwise, α is *essential*. We define the *breadth* $b(F)$ of F as the maximal number of mutually disjoint, mutually non-parallel essential loops in F.

Let F be a closed 2-manifold embedded in S^3 with $b(F) > 0$. We say that F is *compressible* in S^3 if there exists a disk D embedded in S^3 such that $D \cap F = \partial D$ and ∂D is essential in F. Such a disk is called a *compressing disk* for F. Then by cutting F along ∂D, and pasting two parallel copies of D to its boundaries, we obtain another closed 2-manifold F' with $b(F') < b(F)$. Such an operation is called a *compression along* D. Conversely, if F' is obtained from F by a compression along D, then there exists a dual arc α with respect to D, that is, α intersects D in one point and $\alpha \cap F' = \partial\alpha$ such that F can be recovered from F' by tubing along α. See Figure 1.



FIGURE 1. Compression along D

Remark 1.7. We remark that if $b(F) = 0$, then F consists of only 2-spheres and by the Alexander's theorem [1], F is unknotted in S^3. On the other hand, we remark that if $b(F) > 0$, then by [2] or [3], F is *compressible* in S^3.

Theorem 1.8. *Any closed 2-manifold embedded in the 3-sphere can be unknotted by twistings.*

Proof. Let F be a closed 2-manifold consisting of n closed surfaces $F_1, \ldots, F_n$ embedded in S^3. We will prove Theorem 1.8 by induction on the breadth $b(F)$.

In the case of $b(F) = 0$, by Remark 1.7, F is unknotted.

Next suppose that when $b(F) < b$, Theorem 1.8 holds, and assume that $b(F) = b$. Then by Remark 1.7, there exists a compressing disk D for F. Let F' be the closed 2-manifold obtained from F by a compression along D. Then there exists an arc α such that α intersects D in one point, $\alpha \cap F' = \partial\alpha$, and F can be obtained from F' by tubing along α. Since $b(F') < b$, by the assumption of induction, F' can be unknotted by twistings. Thus there exists a sequence of twistings

$$(S^3, F') \xrightarrow{C_1} (S^3, F'(1)) \xrightarrow{C_2} \cdots \xrightarrow{C_m} (S^3, F'(m)),$$

where $F'(m)$ is unknotted. In each stage, we may assume that $C_i \cap \alpha = \emptyset$ for $i = 1, \ldots, m$. Therefore, this sequence extends to a sequence of twistings

$$(S^3, F) \xrightarrow{\mathrm{C_1}} (S^3, F(1)) \xrightarrow{\mathrm{C_2}} \cdots \xrightarrow{\mathrm{C_m}} (S^3, F(m)).$$

Let R be the closure of a connected component of $S^3 - F'$ which contains α, and put $\partial R = F'_1 \cup \cdots \cup F'_k$, where $F'_1, \ldots, F'_k$ are connected components of $F'(m)$. Since $F'(m)$ is unknotted, $F'_1 \cup \cdots \cup F'_k$ bounds k handlebodies $V_1, \ldots, V_k$ in $S^3 - intR$, and $V_1 \cup \cdots \cup V_k$ is unknotted in S^3, namely, $V_1 \cup \cdots \cup V_k$ is ambient isotopic to a regular neighborhood of a plane graph G on the 2-sphere S. Then by crossing changes on α and crossing changes between α and V_i, α can be unknotted, that is, α is isotopic to an arc on S. Since these crossing changes are obtained by twistings, there is a sequence of twistings

$$(S^3, F'(m)) \xrightarrow{\mathrm{C_{m+1}}} (S^3, F'(m+1)) \xrightarrow{\mathrm{C_{m+2}}} \cdots \xrightarrow{\mathrm{C_{m+l}}} (S^3, F'(m+l)),$$

where $F'(m), F'(m+1), F'(m+2), \ldots, F'(m+l)$ are equivalent and $C_{m+i} \cap \alpha = \emptyset$ for $i = 1, \ldots, l$. Therefore, this sequence extends to a sequence of twistings

$$(S^3, F(m)) \xrightarrow{\mathrm{C_{m+1}}} (S^3, F(m+1)) \xrightarrow{\mathrm{C_{m+2}}} \cdots \xrightarrow{\mathrm{C_{m+l}}} (S^3, F(m+l)).$$

Hence, by tubing F' along α, we obtain a sequence of twistings

$$(S^3, F) \xrightarrow{\mathrm{C_1}} (S^3, F(1)) \xrightarrow{\mathrm{C_2}} \cdots \xrightarrow{\mathrm{C_m}} (S^3, F(m))$$
$$\xrightarrow{\mathrm{C_{m+1}}} (S^3, F(m+1)) \xrightarrow{\mathrm{C_{m+2}}} \cdots \xrightarrow{\mathrm{C_{m+l}}} (S^3, F(m+l)),$$

where $F(m+l)$ is unknotted. □

By the Waldhausen's theorem [9], any unknotted closed surface in S^3 is unique up to isotopy. Therefore, by Theorem 1.8, we have the following.

Corollary 1.9. *Any two closed 2-manifolds $F = F_1 \cup \cdots \cup F_n$ and $F' = F'_1 \cup \cdots \cup F'_n$ in the 3-sphere can be mutually related by twistings if $n \leq 2$ and F is homeomorphic to F'.*

Example 1.10. *We recall an example of closed surface H of genus 2 given by Homma* [3]*, see also* [6, 4.1 Theorem] *as shown in Figure 2. The surface H separates S^3 into two components W_1 and W_2, where W_1 is homeomorphic to the exterior of the 4-crossing Handcuff graph 4_1 in the table of* [4]*, and W_2 is a boundary connected sum of two trefoil knot exteriors. It is remarkable that H is incompressible in W_1, whereas H has only one compressing disk D in W_2 up to isotopy by* [8], [7].

In spite of Theorem 1.8, there is a following phenomenon.

Theorem 1.11. *The Homma's surface H cannot be unknotted by twistings in W_1.*

Proof. The key of proof is that any twistings in W_1 can take effect only crossing changes on the tube at D. Then one can show that the Handcuff graph 4_1 cannot be unknotted by crossing changes only on α.

Let C be a loop in W_1 which is trivial in S^3. Let D be a compressing disk for H in W_2 which defines the boundary connected sum of two trefoil knot exteriors as shown in Figure 2. By compressing H along D, we obtain two tori T_1 and T_2 which bound two trefoil knot exteriors E_1 and E_2 in W_2. Let α be an arc in W_2 which intersects D in one point and connects T_1 and T_2 such that H is obtained from $T_1 \cup T_2$ by tubing along α. Since C is the trivial knot in S^3, both of T_1



FIGURE 2. The Homma's closed surface



FIGURE 3. The 4-crossing Handcuff graph 4_1

and T_2 are compressible in $S^3 - int(E_1 \cup E_2) - C$. Hence C bounds a disk Δ in $S^3 - int(E_1 \cup E_2)$. Then we may assume that α intersects Δ transversely and conclude that any twisting along C takes effect only on α.

Let $K_1 \cup \alpha \cup K_2$ be the Handcuff graph 4_1, whose exterior is homeomorphic to W_1. We take a double branched cover of S^3 along the trivial link $K_1 \cup K_2$ as follows. Let D_i be a disk bounded by K_i which intersects α in one point $(i = 1, 2)$. We cut open S^3 along $D_1 \cup D_2$ and take a copy of it. Those 3-manifolds are both homeomorphic to $S^2 \times I$ and whose boundary consists of 2-spheres $D_1^+ \cup D_1^-$, $D_2^+ \cup D_2^-$, $D'^+_1 \cup D'^-_1$, $D'^+_2 \cup D'^-_2$. Then by gluing D_1^+ and D'^-_1, D_1^- and D'^+_1, D_2^+ and D'^-_2, D_2^- and D'^+_2 we obtain $S^2 \times S^1$ and a knot $\tilde{\alpha}$ obtained from α and α' as shown in Figure 4. We note that $[\tilde{\alpha}] = 3[\gamma]$ in $H_1(S^2 \times S^1; \mathbb{Z}) \cong \mathbb{Z}$, where γ is a generator of $H_1(S^2 \times S^1; \mathbb{Z})$.

Suppose that $K_1 \cup \alpha \cup K_2$ is unknotted by crossing changes on α. Then the homology class $[\tilde{\alpha}]$ in $H_1(S^2 \times S^1; \mathbb{Z})$ does not change by the crossing changes, and we have $[\tilde{\alpha}] = 3[\gamma]$. However, since $K_1 \cup \alpha \cup K_2$ is trivial, we have $[\tilde{\alpha}] = [\gamma]$. This is a contradiction. □



FIGURE 4. The double branched cover of S^3 along $K_1 \cup K_2$

By the Fox's re-embedding theorem [2], there exists a re-embedding of W_2 into S^3 such that W_2 is unknotted. However, this Fox's re-embedding cannot be obtained by twistings.

Corollary 1.12. *There exists a 3-submanifold of S^3 which cannot be unknotted by twistings.*

Proof. Take W_2 as a 3-submanifold of S^3. □

However, we remark that by [5, Theorem 1.6], there exists a null-homologous link L in W_1, which is reflexive in S^3, such that a handlebody can be obtained from W_1 by a $1/\mathbb{Z}$-Dehn surgery along L, that is, $[L] = 0$ in $H_1(W_1; \mathbb{Z})$ and there exists a surgery slope $1/n_i$ for each component L_i of L such that a pair of S^3 and a handlebody is obtained from (S^3, W_1) by a Dehn surgery along L. Therefore, the Fox's re-embedding can be replaced with a Dehn surgery along a link. At the time of writing of [5], it was unknown whether this Dehn surgery can be replaced with twistings. Corollary 1.12 shows that it is not always true.

REFERENCES

1. J. W. Alexander, *On the subdivision of a 3-space by a polyhedron*, Proc. Nat. Acad. Sci. USA **10** (1924), 6–8.
2. R. H. Fox, *On the imbedding of polyhedra in 3-space*, Ann. of Math. **49** (1948), 462–470.
3. T. Homma, *On the existence of unknotted polygons on 2-manifolds in E^3*, Osaka Math. J. **6** (1954), 129–134.
4. A. Ishii, K. Kishimoto, H. Moriuchi and M. Suzuki, *A table of genus two handlebody-knots up to six crossings*, J. Knot Theory Ramifications **21**, 1250035 (2012).
5. M. Ozawa, K. Shimokawa, *Dehn surgery and Seifert surface systems*, to appear in Ann. Scuola Norm-Sci.
6. S. Suzuki, *On a complexity of a surface in 3-sphere*, Osaka J. Math. **11** (1974), 113–127.
7. S. Suzuki, *On surfaces in* 3*-sphere: prime decomposition*, Hokkaido Math. J. **4** (1975), 179–195.
8. Y. Tsukui, *On surfaces in* 3*-space*, Yokohama Math. J. **18** (1970), 93–104.
9. F. Waldhausen, *Heegaard-Zerlegungen der* 3*-Sphäre*, Topology **7** (1968), 195–203.

DEPARTMENT OF NATURAL SCIENCES, FACULTY OF ARTS AND SCIENCES, KOMAZAWA UNIVERSITY, 1-23-1 KOMAZAWA, SETAGAYA-KU, TOKYO, 154-8525, JAPAN
E-mail address: w3c@komazawa-u.ac.jp